

Todd Rowe


Argemone pleicantha - A Proving of Crested Prickle Poppy

Extrait du livre

[Argemone pleicantha - A Proving of Crested Prickle Poppy](#)

de [Todd Rowe](#)

Éditeur : American Medical College of Homeopathy Publishing, Inc.



<http://www.editions-narayana.fr/b2005>

Sur notre [librairie en ligne](#) vous trouverez un grand choix de livres d'homéopathie en français, anglais et allemand.

Reproduction des extraits strictement interdite.

Narayana Verlag GmbH, Blumenplatz 2, D-79400 Kandern, Allemagne

Tel. +33 9 7044 6488

Email info@editions-narayana.fr

<http://www.editions-narayana.fr>



Crested Prickle Poppy

Table of Contents

Introduction	Page 1
Natural History of Crested Prickle Poppy	Page 3
Symptom Listing and suggested Repertory additions	Page 5
Ideas and Themes	Page 17
Comparison to Argemone mexicana	Page 29
Crested Prickle Poppy Raw Data	Page 33
Discussion	Page 55
Bibliography	Page 59
Appendix A:	Page 61
Comparison to other Papaveraceae Remedies	
Appendix B:	Page 71
Comparison to Flower Essence Proving of Crested Prickle Poppy	

Introduction




Proving Methodology

A proving of Crested Prickle Poppy (*Argemone pleiakantha*) was performed in Phoenix Arizona in the fall of 2000. The proving was conducted by the Society for the Establishment of Research in Classical Homeopathy (SERCH), which is a division of the Desert Institute of Classical Homeopathy. The proving consisted of thirteen subjects who began taking *Argemone pleiakantha* 30C on September 1, 2000. The proving was double blinded, which means that neither the provers nor the supervisors knew the identity of the remedy until the proving was completed.

The instructions were to take a single daily dosage for a maximum of three days, but to stop after any symptoms became apparent. Participants journaled their symptoms over the space of the next month. Informed consent was obtained from each participant. Symptoms were reviewed in an exit group meeting one month later. This session was videotaped. Journals were reviewed separately. There were eight proving supervisors who followed the provers closely over the months following the start of the proving. The proving supervisors reported directly to the proving coordinator.

This was a partial proving. There were five male participants and eight female participants. Their ages ranged from 28-59 years.

The specimen was obtained from Eileen Nauman DHM (UK). She prepared the substance on April 5, 1995 from a plant in the Sonoran Desert north of Phoenix, Arizona. The remedy was prepared from a mother tincture of the flower.



Todd Rowe

[Argemone pleicantha - A Proving of
Crested Prickle Poppy](#)

86 pages, reliure spirales
publication 2007



Plus de livres sur homéopathie, les médecines naturelles et un style de vie plus sain
www.editions-narayana.fr