


B. Jain Surgery (Paper II)-Ent, Eye & Dental

Extrait du livre

[Surgery \(Paper II\)-Ent, Eye & Dental](#)

de [B. Jain](#)

Éditeur : B. Jain



<http://www.editions-narayana.fr/b5590>

Sur notre [librairie en ligne](#) vous trouverez un grand choix de livres d'homéopathie en français, anglais et allemand.

Reproduction des extraits strictement interdite.

Narayana Verlag GmbH, Blumenplatz 2, D-79400 Kandern, Allemagne

Tel. +33 9 7044 6488

Email info@editions-narayana.fr


<http://www.editions-narayana.fr>



Treatment :

- (1) If Pterygium is stationary for atrophic stage and not much disfiguring, no treatment is necessary.
- (2) In progressive pterygium, the surgery is required.

(ii) Myopia : It is the error of refraction in which the parallel rays of light from infinity come to a focus in front of retina, when accommodation is at rest.

**Types of Myopia :**

- (1) Axial myopia
- (2) Curvature myopia
- (3) Index myopia

Clinical types :

- (i) Congenital myopia
- (ii) Simple myopia
- (iii) Pathological myopia

Ophthalmoscopic findings :

- (a) Myopic crescent.
- (b) Chorio-retinal atrophy.
- (c) Atrophic patches in the macula with sclerosis of choroidal vessels.
- (d) Ectasia of sclera at the posterior pole known as *posterior staphyloma*.

Symptoms of Myopia :

- Reduced visual acuity.
- But near objects seen clearly.

- The patient may complain of seeing black spots floating in front of eyes due to vitreous opacities.

Treatment : By concave lens of suitable power.

(iii) Leucoplakia : Leucoplakia is the appearance of white patch usually on the cords and may look like a small prominent patch or a nodular plaque.

The lesion may be single or multiple. Leucoplakia is generally associated with hyperkeratosis.

- Causes**
- Idiopathic
 - Tobacco chewing

Prognosis - Leucoplakia is premalignant dysplasia.


Treatment - Microsurgical excision.

(iv) Pyorrhea : Pyorrhea is an inflammatory infective condition of the gums and the membranes covering the teeth, in which there is a discharge of pus from teeth sockets or a collection of pus between the teeth and the gum margins.

Causes - The most frequent single cause of gingivitis (Pyorrhea) is poor dental hygiene with a buildup of dental plaque around the base of the teeth. Other localised causes include tartar (calcified plaque), food traps between teeth and ill-fitting dental plates, caps or crowns. The gum inflammation that often develops during puberty or pregnancy is thought to be aggravated by hormonal factors. Gingivitis may also be a symptom of general illness (for example, in Diabetes Mellitus and Leukemia) and malnutrition (especially in scurvy and pellagra-Vitamin B₃ deficiency).

Complications :

- (1) Bleeding from gums.
- (2) An perverted taste due to pus in mouth.
- (3) Recession of gums and loosening of teeth.



B. Jain

Surgery (Paper II)-Ent, Eye & Dental

192 pages, broché
publication 2002



acheter maintenant

Plus de livres sur homéopathie, les médecines naturelles et un style de vie plus sain

www.editions-narayana.fr