


Samuel Hahnemann

The Chronic Diseases

Extrait du livre

[The Chronic Diseases](#)
de [Samuel Hahnemann](#)

Éditeur : B. Jain



<http://www.editions-narayana.fr/b1359>

Sur notre [librairie en ligne](#) vous trouverez un grand choix de livres d'homéopathie en français, anglais et allemand.

Reproduction des extraits strictement interdite.

Narayana Verlag GmbH, Blumenplatz 2, D-79400 Kandern, Allemagne

Tel. +33 9 7044 6488

Email info@editions-narayana.fr

<http://www.editions-narayana.fr>



SYCOSIS

First, then, concerning sycosis, as being that miasma which has produced by far the fewest chronic diseases, and has only been dominant from time to time. This *fig-wart disease*, which in later times, especially during the French war, in the years 1809-1814, was so widely spread, but which has since showed itself more and more rarely, was treated, almost always, in an inefficient and injurious manner, internally with Mercury, because it was considered homogeneous with the venereal chancre-disease; but the excrescences on the genitals were treated by Allopathic physicians always in the most violent external way by cauterizing, burning and cutting, or by ligatures. These excrescences usually first manifest themselves on the genitals, and appear usually, but not always, attended with a sort of gonorrhoea [Usually in gonorrhoea of this kind, the discharge is from the beginning thickish, like pus; micturition is less difficult, but the body of the penis swollen somewhat hard; the penis is also in some cases covered on the back with glandular tubercles, and very painful to the touch.] from the urethra, several days or several weeks, even many weeks after infection through coition; more rarely they appear dry and like warts, more frequently sort, spongy, emitting a specifically fetid fluid (sweetish and almost like herring-brine), bleeding easily, and in the form of a coxcomb or a cauliflower (*brassica botrytes*). These, with males, sprout forth on the glans and on, or below, the prepuce, but with women, on the parts surrounding the pudenda; and the pudenda themselves, which are then swollen, are covered often by a great number of them. When these are violently removed, the natural, proximate effect is, that they will usually come forth again, usually to be subjected again, in vain, to a similar, painful, cruel treatment. But even if they could be rooted out in this way, it would merely have the consequence, that the fig-wart disease, after


having been deprived of the local symptom which acts vicariously for the internal ailment, would appear [The miasm of the other common gonorrhoea seems not to penetrate the whole organism, but only to locally stimulate the urinary organs. They yield either to a dose of one drop of fresh parsley-juice, when this is indicated by a frequent urgency to urinate, or a small dose of Cannabis, of Cantharides, or of the Copaiva balm, according to their different constitution and the other ailments attending it. These should, however, be always used in the higher and highest dynamizations (potencies), unless a Psora, slumbering in the body of the patient, has been developed by means of a strongly affecting, irritating or weakening treatment by Allopathic physicians. In such a case frequently secondary gonorrhoeas remain, which can only be cured by an anti-psoric treatment.] in other and much worse ways, in secondary ailments; for the fig-wart miasm, which rules in the whole organism, has been in no way diminished, either by the external destruction of the above-mentioned excrescences, or by the Mercury which has been used internally and which is in no way appropriate to sycosis. Besides the undermining of the general health by Mercury, which in this disease can only do injury, and which is given mostly in very large doses and in the most active preparations, similar excrescences then break out in other parts of the body, either whitish, spongy, sensitive, flat elevations, in the cavity of the mouth, on the tongue, the palate and the lips, or as large, raised, brown and dry tubercles in the axillae, on the neck, on the scalp, etc., or there arise other ailments of the body of which I shall only mention the contraction of the tendons of the flexor muscles, especially of the fingers.

The gonorrhoea dependent on the fig-wart miasma, as well as the above-mentioned excrescences (*i.e.*, the whole sycosis), are cured most surely and most thoroughly through the internal use of Thuja, [Materia Medica Pura, Part V.] which, in this case, is Homoeopathic, in a dose of a few pellets as large as poppy seeds, moistened with the dilution potentized to the decillionth [If further doses of Thuja are required, they are used most efficiently from other potencies (viii., vi., v., ii.), a change of the modification of the remedy, which facilitates and strengthens its ability of affecting the vital force,] degree, and when these have exhausted their action after fifteen, twenty, thirty, forty days, alternating with just as small a dose of Nitric acid, diluted to the decillionth degree, which must be allowed to act as long a time, in order to remove the gonorrhoea and the excrescences; *i.e.*, the whole sycosis. It is not necessary to use any

external application, except in *the most inveterate and difficult* cases, when the larger fig-warts may be moistened every day with the mild, pure juice pressed from the green leaves of Thuja, mixed with an equal quantity of Alcohol.

But if the patient was at the same time affected with another chronic ailment, as is usual after the violent treatment of fig-warts by Allopathic physicians, then we often find developed Psora [This Psora is hardly ever found in its developed state (and thus capable of entering into complication with other miasmata) with young people who have just been infected and seized by the fig-wart disease, and who have not had to pass through the usual Mercurial treatment, which never runs its course without the most violent assaults on the constitution; by this pernicious derangement of the whole organism, the Psora, even if slumbering ever so soundly, will be awakened, if as is often the case, it was present within.] complicated with sycosis, when the Psora, as is often the case, was latent before in the patient. At times, when a badly treated case of venereal chancre disease had preceded, both these miasmata are conjoined in a three-fold complication with syphilis. Then it is necessary first to come to the assistance of the most afflicted part, the *Psora*, with the specific anti-psoric remedies given below, and then to make use of the remedies for sycosis, before the proper dose of the best preparation of Mercury, as will be described below, is given against the syphilis; the same alternating treatment may be continued, until a complete cure is effected. Only, each one of these three kinds of medicine must be given the proper time to complete its action.

In this reliable cure of sycosis from within, no external remedy (except the juice of Thuja in inveterate bad cases) must be applied or laid on the fig-warts, only clean, dry lint, if they are of the moist variety.



Samuel Hahnemann

[The Chronic Diseases](#)

Their Peculiar Nature and Their
Homoeopathic Cure

218 pages, broché
publication 2015



Plus de livres sur homéopathie, les médecines naturelles et un style de vie plus sain
www.editions-narayana.fr